



IN THE CLAIMS

Kindly amend the claims as shown on the attached sheets.

We claim:

1. (original) A process for preparing indenenes of the formula (II) from indenenes of the formula (I) by reaction with compounds of the formula (III)



15 where

$X^1$  is chlorine;

20  $Y^1$  are identical or different and are selected independently from the group consisting of  $C_1$ - $C_{40}$  groups, e.g.  $C_1$ - $C_{25}$ -alkyl,  $C_2$ - $C_{25}$ -alkenyl,  $C_2$ - $C_{25}$ -alkynyl,  $C_3$ - $C_{15}$ -alkylalkenyl,  $C_3$ - $C_{15}$ -alkylalkynyl,  $C_6$ - $C_{24}$ -aryl,  $C_4$ - $C_{24}$ -heteroaryl,  $C_5$ - $C_{24}$ -alkylheteroaryl,  $C_7$ - $C_{30}$ -arylalkyl,  $C_7$ - $C_{30}$ -alkylaryl,  $C_1$ - $C_{12}$ -alkoxy,  $C_6$ - $C_{24}$ -aryloxy, fluorinated  $C_1$ - $C_{25}$ -alkyl, fluorinated  $C_6$ - $C_{24}$ -aryl, fluorinated  $C_7$ - $C_{30}$ -arylalkyl, fluorinated  $C_7$ - $C_{30}$ -alkylaryl, and the fluorine atom and heteroatom-containing groups, e.g. boron-, silicon-, nitrogen-, oxygen- or sulfur-containing groups, which may bear one or more substituents, where a plurality of groups  $Y^1$  may also together form a cyclic aliphatic or aromatic ring system which may in turn be substituted and may contain heteroatoms;

30  $Y^2$  are identical or different and are selected independently from the group consisting of  $C_1$ - $C_{40}$  groups, e.g.  $C_1$ - $C_{25}$ -alkyl,  $C_2$ - $C_{25}$ -alkenyl,  $C_2$ - $C_{25}$ -alkynyl,  $C_3$ - $C_{15}$ -alkylalkenyl,  $C_3$ - $C_{15}$ -alkylalkynyl,  $C_6$ - $C_{24}$ -aryl,  $C_4$ - $C_{24}$ -heteroaryl,  $C_5$ - $C_{24}$ -alkylheteroaryl,  $C_7$ - $C_{30}$ -arylalkyl,  $C_7$ - $C_{30}$ -alkylaryl,  $C_1$ - $C_{12}$ -alkoxy,  $C_6$ - $C_{24}$ -aryloxy, fluorinated  $C_1$ - $C_{25}$ -alkyl, fluorinated  $C_6$ - $C_{24}$ -aryl, fluorinated  $C_7$ - $C_{30}$ -arylalkyl, fluorinated  $C_7$ - $C_{30}$ -alkylaryl, and heteroatom-containing groups, e.g. boron-, silicon-, nitrogen-, oxygen- or sulfur-containing groups, which may bear one or more substituents, where a plurality of groups  $Y^2$  may also together form a cyclic aliphatic or aromatic ring system which may in turn be substituted and may contain heteroatoms;

40

5  $R^1$  are identical or different and are selected independently from the group consisting of linear, branched or cyclic aliphatic hydrocarbon groups, e.g.  $C_1$ - $C_{25}$ -alkyl which may in turn bear a variety of substituents, and groups bound via an aliphatic group to the indenyl skeleton, e.g.  $C_3$ - $C_{15}$ -alkenylalkyl,  $C_3$ - $C_{15}$ -alkynylalkyl,  $C_5$ - $C_{24}$ -heteroarylalkyl,  $C_7$ - $C_{30}$ -arylalkyl,  $C_2$ - $C_{30}$ -alkyloxyalkyl,  $C_7$ - $C_{30}$ -aryloxyalkyl,  $C_8$ - $C_{30}$ -alkylarylalkyl, and other heteroatom-containing groups which are bound via an aliphatic group to the indenyl skeleton, e.g. boron-, silicon-, nitrogen-, oxygen- or sulfur-containing groups, and may bear one or more substituents;

10  $M^1$  is an element of group 1, 2, 12, 13 or 14 of the Periodic Table of the Elements;

15  $X^2$  are identical or different and are selected independently from the group consisting of halogen atoms, the hydroxy group, alkoxy groups, aryloxy groups, mesylate, tosylate and triflate;

$m$  is an integer from 0 to 3;

$n$  is an integer from 1 to 4;

$p$  is an integer from 0 to 4;

$q$  is an integer from 1 to 4;

20  $r$  is 1, 2 or 3, and

$t$  is 0, 1 or 2, where  $r + t$  corresponds to the oxidation number of  $M^1$ ;

wherein the indenenes of the formula (I) are reacted with appropriate aliphatic organometallic compounds of the formula (III) in the presence of a transition metal catalyst.

25 2. (origina) A process as claimed in claim 1, wherein

$M^1$  is Li, Mg, B or Zn, and

$X^2$  are identical or different and are selected independently from the group consisting of halogen atoms, the hydroxy group, alkoxy groups and aryloxy groups.

30 3. (currently amended) A process as claimed in claim 1 or 2, wherein at least one transition metal

catalysts selected from the group consisting of nickel(II) acetylacetonate, [1,2-bis-(diphenylphosphino)ethane]nickel(II) chloride, [1,3-

bis(diphenylphosphino)propane]nickel(II)chloride, [1,1'-bis(diphenylphos-

35 phino)ferrocene]nickel(II) chloride, bis(tributylphosphine)nickel(II) bromide,

bis(tributylphosphine)nickel(II) chloride, bis(triphenylphosphine)nickel(II) chloride,

bis(triphenylphosphine)dicarbonylnickel(0), [1,2-bis(dimethylphosphino)ethane]nickel(II)


chloride, bis(triethylphosphine)nickel(II) chloride, bis(triphenylphosphine)palladium(II)

chloride, tetrakis(triphenylphosphine)palladium(0), [1,2-bis(diphenylphosphino)-

40

ethane]palladium(II) chloride and the [1,1'-bis(diphenylphosphino)ferrocene]nickel(II) chloride-methylene chloride complex is used.

4. (currently amended) A process as claimed in claim 1 any of claims 1 to 3, wherein the transition metal catalyst used is [1,3-bis(diphenylphosphino)propane]nickel(II) chloride.
5. (currently amended) A process as claimed in claim 1 any of claims 1-4, wherein the transtransition metal catalyst is added in an amount of from 0.01 to 5 mol%, based on chloroindene of the formula (I) used.
- 10 6. (currently amended) A process as claimed in claim 1 any of claims 1-5, wherein the chloroindenes of the formula (I) which are used are selected from the groups consisting of:  
 4-chloro-1-indene; 5-chloro-1-indene; 6-chloro-1-indene; 7-chloro-1-indene; 2-methyl-4-chloro-1-indene; 2,7-dimethyl-4-chloro-1-indene; 2,4-dimethyl-7-chloro-1-indene; 2-methyl-5-chloro-1-indene; 2-methyl-6-chloro-1-indene; 2-methyl-7-chloro-1-indene; 2-ethyl-4-chloro-1-indene; 2-ethyl-5-chloro-1-indene; 2-ethyl-6-chloro-1-indene; 2-ethyl-7-chloro-1-indene; 2-propyl-4-chloro-1-indene; 2-propyl-5-chloro-1-indene; 2-propyl-6-chloro-1-indene; 2-propyl-7-chloro-1-indene; 2-i-propyl-4-chloro-1-indene; 2-i-propyl-5-chloro-1-indene; 2-i-propyl-6-chloro-1-indene; 2-i-propyl-7-chloro-1-indene; 2-butyl-4-chloro-1-indene; 2-butyl-5-chloro-1-indene; 2-butyl-6-chloro-1-indene; 2-butyl-7-chloro-1-indene; 2-s-butyl-4-chloro-1-indene; 2-s-butyl-5-chloro-1-indene; 2-s-butyl-6-chloro-1-indene; 2-s-butyl-7-chloro-1-indene; 2-t-butyl-4-chloro-1-indene; 2-t-butyl-5-chloro-1-indene; 2-t-butyl-6-chloroindene; 2-t-butyl-7-chloroindene.
- 20 7. (original) An indene of the formula (IIa) or (IIb),



30

where

$R^2$  is  $C_1$ - $C_{10}$ -alkyl and

$R^3$  is a monocyclic or polycyclic  $C_5$ - $C_{15}$ -alkyl group or a  $CH_2R^4$  group,

where

35  $R^4$  is a  $C_6$ - $C_{14}$ -aryl group, a  $C_7$ - $C_{15}$ -alkylaryl group or a monocyclic or polycyclic  $C_5$ - $C_{15}$ -alkyl group.

8. (currently amended) The use of indenenes of the formula (II) obtainable as set forth in any of claims 1 to 6 or indenenes of the formula (IIa) or (IIb) as claimed in claim 9 as intermediates in the synthesis of active compounds or A process for the synthesis of metallocene compounds, said process comprising synthesizing the compounds from indenenes of the formula (II) obtained as set forth in claim 1.

9. (original) An ansa-bisindenylmetallocene prepared from at least one indene of the formula (IIa) or (IIb) as claimed in claim 7, wherein the two indenyl ligands of the metallocene have different substituents in the 2 position.
10. (currently amended) The use of A process for preparing olefins, said process comprising polymerizing said olefins in the presence of at least one ansa-bisindenylmetallocene prepared from at least one indene of the formula (II) as set forth in ~~any of claims 1 to 6~~ claim 1 ~~or an ansa-bisindenylmetallocene as claimed in claim 9 for the polymerization of olefins.~~

15

20

25

30

35

40